

TPI-MINN-31-97
HEP-97-1617
hep-ph/9712323

HQET AND SEMILEPTONIC FORM FACTORS

ARKADY VAINSHTEIN

*Theoretical Physics Institute, University of Minnesota
and*

Budker Institute of Nuclear Physics, Novosibirsk

*Talk at the Seventh International Symposium On Heavy Flavor Physics,
July 7-11, 1997, Santa Barbara;
to appear in the Proceedings*

Abstract

Theoretical approaches to form factors of semileptonic decays are discussed in application to $B \rightarrow D + l + \bar{\nu}_l$ decay.

HQET AND SEMILEPTONIC FORM FACTORS

ARKADY VAINSHTEIN

Theoretical Physics Institute, University of Minnesota

and

Budker Institute of Nuclear Physics, Novosibirsk

Theoretical approaches to form factors of semileptonic decays are discussed in application to $B \rightarrow D + l + \bar{\nu}_l$ decay.

1 Outline

Heavy Quark Effective Theory (HQET)¹ is a powerful approach to the processes involving heavy flavor hadrons and, in particular, to the study of form factors of semileptonic decays (see reviews²). From a field-theoretical standpoint, the HQET is a particular example of construction based on a more general notion of Wilson Operator Product Expansion (OPE) or Wilson effective action³, to which I mainly refer.

The outline of the talk is as follows:

- Heavy flavor decays as short distance probes. Total semileptonic widths
- From inclusive to exclusive: bounds for $B \rightarrow D^*$ leading form factor
- Subleading $B \rightarrow D^*$ form factors
- Some recent developments
- Conclusions

2 Heavy Flavor Decays as Short Distance Probes

The quark picture of semileptonic B meson decay $B \rightarrow D + e + \bar{\nu}_e$ is presented by Fig. 1. (Better to put $\bar{B}$ but I will systematically neglect by bar on B .) The specific feature of heavy flavor states is that they contain a heavy quark Q which could be viewed as an almost static object (in the rest frame of hadron). Thus these states can be considered as ground states of light flavor QCD in the presence of static source of gluon field. A relevant theoretical parameter is

$$\frac{\Lambda_{QCD}}{m_Q} \ll 1. \quad (1)$$




Figure 1: Semileptonic $B \rightarrow D + e + \bar{\nu}_e$ decay

An immediate consequence is so called spin-flavor symmetry, i.e., an independence on heavy quark spin and flavor in the large mass limit where the relation (1) holds.

However, the characteristic size of heavy flavor hadron is of order $1/\Lambda_{QCD}$ and the structure of heavy flavor states as well as that of the QCD vacuum is governed by nonperturbative dynamics in strong coupling regime. To see a simple underlying quark-gluon structure at short distances, where the strong interaction becomes weaker one needs a short distance probe. Heavy flavor decays, particularly the ones inclusive over final hadrons, provide such a probe. This probe is similar to the famous ratio R for the inclusive cross section of hadron production with a substitution of QCD vacuum by a heavy flavor hadron. This comparison is illustrated by Figs. 2 and 3. Fig. 2 is for the process of e^+e^- inclusive annihilation into hadrons, its cross section is given by

$$R = \frac{\sigma(e^+e^- \rightarrow \text{hadrons})}{\sigma(e^+e^- \rightarrow \mu^+\mu^-)} = N_{\text{colors}} \sum_q Q_q^2 [1 + O[\alpha_S(W)] + \dots], \quad (2)$$

in the limit of high total energy W , i.e., when $W \gg \Lambda_{QCD}$. Fig. 3 presents semileptonic decays of B meson inclusive over nonstrange hadrons in the final




Figure 2: e^+e^- annihilation into hadrons (inclusive)

state ($b \rightarrow u$ transition in terms of quarks),

$$\Gamma_{SL}^{ub} = \sum_{X_u} \Gamma(B \rightarrow X_u e \bar{\nu}) = \frac{G_F^2}{192\pi^3} m_b^5 |V_{ub}|^2 [1 + O[\alpha_S(m_b)] + \dots] \quad (3)$$

in the limit of large mass of b quark, $m_b \gg \Lambda_{QCD}$.

In both examples the leading term is given by partonic description of the process where only hard momenta count and an influence of soft background of quarks and gluons in the vacuum and in B meson is strongly suppressed. Theoretical basis of quark-hadron duality is not trivial and based on two notions:

- (i) Wilson Operator Product Expansion³. The expansion takes place at large *Euclidean* momenta where no hadron can be produced. The OPE means a factorization of short and large distances in Euclidean domain.
- (ii) Dispersion relations. Based on causality dispersion relations express certain integrals over observable cross sections (in *Minkowski* domain) as correlators at Euclidean momenta.

The construction described allows for calculation of corrections to the leading partonic term, both perturbative, $[\alpha_S(m_b)]^n$, and nonperturbative, $[\Lambda_{QCD}/m_b]^k$. For the semileptonic total width (3) it results in

$$\Gamma_{SL}^{ub} = \frac{G_F^2 m_b^5 |V_{ub}|^2}{192\pi^3} \left[1 - \frac{2\alpha_S}{3\pi} - \frac{\mu_\pi^2}{2m_b^2} - \frac{3\mu_G^2}{2m_b^2} \right] \quad (4)$$




Figure 3: Inclusive width of $B \rightarrow X_u e \bar{\nu}$ decays

where power corrections⁴ are defined by parameters μ_π^2 and μ_G^2 . The first one, μ_π^2 , has the meaning of mean value for momentum square of b quark inside of B meson,

$$\mu_\pi^2 = \langle B | \bar{b} \pi^2 b | B \rangle \approx 0.6 \pm 0.1 \text{ GeV}^2 . \quad (5)$$

The value given is due to theoretical estimate⁵, the direct determination is not done yet. The second parameter, μ_G^2 , measures chromomagnetic interaction with the spin of heavy quark, so its value is fixed by spin splitting in B , B^* system,

$$\mu_G^2 = \langle B | \bar{b} \frac{i}{2} \sigma^{\mu\nu} G_{\mu\nu} b | B \rangle = \langle \vec{\sigma} \vec{B} \rangle = \frac{3}{4} (M_{B^*}^2 - M_B^2) \approx 0.36 \text{ GeV}^2 . \quad (6)$$

3 From Inclusive to Exclusive

The decays $B \rightarrow D^* l \bar{\nu}_l$ and $B \rightarrow D l \bar{\nu}_l$ produce about 2/3 of the total semileptonic width. The suppression of production of higher states can be understood as due to a closeness to the point of zero recoil. In other words, hadrons in the final state are slow moving. The point of zero recoil, where the spatial momentum $\vec{q} = 0$, is a special one in the case of heavy quark in the final state. The quark-hadron duality takes place at this point in spite of small momentum transfer. It is a remarkable case where OPE works up to large distances (in

case of light quarks a similar phenomena shows up in the axial current anomaly and in the certain term of OPE for the product of two axial currents).

The theoretical knowledge of the amplitude of $B \rightarrow D^* l \bar{\nu}_l$ decay at zero recoil is used, in particular, for an extraction of $|V_{cb}|$ from the data. The crucial for this extraction question is about a theoretical accuracy of HQET predictions.

Let us start with kinematics of the decay. There are four invariant form factors describing the transition²,

$$\begin{aligned} \langle D^* | \vec{V} - \vec{A} | B \rangle &= \sqrt{m_B m_{D^*}} h_{A_1}(w) [-(w+1)\vec{e}_{D^*} + \\ &w R_1(w) i \vec{v}_{D^*} \times \vec{e}_{D^*} + w R_3(w) \vec{v}_{D^*} (\vec{v}_{D^*} \cdot \vec{e}_{D^*})] , \end{aligned} \quad (7)$$

$$\begin{aligned} \langle D^* | V_0 - A_0 | B \rangle &= \sqrt{m_B m_{D^*}} h_{A_1}(w) (\vec{v}_{D^*} \cdot \vec{e}_{D^*}) [-(w+1) + \\ &R_2(w) \frac{m_B}{m_{D^*}} - R_3(w) \left(\frac{m_B}{m_{D^*}} - 1 \right)] , \end{aligned} \quad (8)$$

where matrix elements of $V - A$ current are written separately for spatial $\vec{V} - \vec{A}$ and timelike $V_0 - A_0$ components and

$$w = \frac{E_{D^*}}{m_{D^*}} = \sqrt{1 + \frac{\vec{q}^2}{m_{D^*}^2}} , \quad \vec{v}_{D^*} = -\frac{\vec{q}}{m_{D^*} w} . \quad (9)$$

The $h_{A_1}(w)$ form factor is the only one contributing at zero recoil. Subleading (in terms of their contribution to the amplitude near zero recoil) form factors are introduced in the form of their ratios R_i to the leading one. In the HQET limit these ratios go 1,

$$R_1(w) = R_2(w) = R_3(w) = 1 \quad (\text{at } m_b, m_c \rightarrow \infty) \quad (10)$$

3.1 Zero Recoil Sum Rule

The amplitude at zero recoil is defined by $h_{A_1}(1)$ as discussed above. To get its value let us write down the sum rule⁶ for *inclusive* transitions $B \rightarrow X_c l \bar{\nu}_l$ at zero recoil, $\vec{q} = 0$,

$$|h_{A_1}(1)|^2 + \sum_{m_{D^*} < m_{X_c} < m_{D^*} + \mu} |h_n(1)|^2 = \xi_A(\mu) - \Delta_{1/m^2} + O\left(\frac{1}{m^3}\right) . \quad (11)$$

The states X_c produced at zero recoil are D^* and its radial excitations and $h_n(1)$ are analogs of $h_{A_1}(1)$ for radial excitations. The production of excitation

is suppressed in the heavy quark limit, $h_n(1) \sim 1/m$ and the sum in the l.h.s. of Eq. (11) is of $1/m^2$ order. At the r.h.s. the quantity $\xi_A(\mu)$ gives a perturbative part and has the form

$$\xi_A(\mu) = 1 + 2 \frac{\alpha_S}{\pi} \left[\frac{m_b + m_c}{m_b - m_c} \ln \frac{m_b}{m_c} - \frac{8}{3} \right] + \frac{2}{3} \frac{\alpha_S}{\pi} \mu^2 \left[\frac{1}{m_c^2} + \frac{1}{m_b^2} + \frac{2}{3m_b m_c} \right] \quad (12)$$

Let me note here that the full calculation of the α_S^2 corrections was recently performed⁷. It turned out to be a small change as compared with the value taken from the BLM procedure. Numerically the result for ξ_A is

$$\xi_A(\mu = 0.5m_c) = 0.99 \pm 0.02 \quad (13)$$

The quantity Δ_{1/m^2} presents nonperturbative effects which are of the second order in $1/m$, Eq. (11) contains no corrections of the first order in $1/m$ in correspondence with Luke theorem¹¹,

$$\Delta_{1/m^2} = \frac{\mu_G^2}{3m_c^2} + \frac{\mu_\pi^2 - \mu_G^2}{4} \left[\frac{1}{m_c^2} + \frac{1}{m_b^2} + \frac{2}{3m_b m_c} \right]. \quad (14)$$

Numerically it is about 0.1 which is not a small effect, particular for the quantity of the second order in $1/m$.

The most uncertain in the use of sum rule (11) for theoretical evaluation of $h_{A_1}(1)$ is the sum over excitation. In the absence of direct experimental information the natural assumption would be to say that the sum is comparable with Δ_{1/m^2} . Indeed, the origin of Δ_{1/m^2} can be traced to the contribution of the highly excited states with the excitation energies of order of m_c while the sum under consideration refers to less excitation energies. If some kind of continuity exists then an assumption

$$\sum_{m_{D^*} < m_{X_c} < m_{D^*} + \mu} |h_n(1)|^2 = (0.5 \pm 0.5) \Delta_{1/m^2} \quad (15)$$

seems reasonable for the range of $\mu \sim 0.5 m_c$.

Summarizing numerical estimates we get (see Ref.⁸ for details)

$$h_{A_1}(1) = 0.91 - 0.013 \frac{\mu_\pi^2 - 0.5 \text{ GeV}^2}{0.1 \text{ GeV}^2} \pm 0.02_{\text{excit}} \pm 0.01_{\text{pert}} \pm 0.025_{1/m^3} \quad (16)$$

where the uncertainties are marked by the their sources. Overall,

$$h_{A_1}(1) = 0.91 \pm 0.06, \quad (17)$$

what leads to the 6% uncertainty in the $|V_{cb}|$ extraction from the exclusive D^* production at zero recoil. The theoretical uncertainty is less by about a factor of two in the determination of $|V_{cb}|$ from inclusive production⁸. The difference is due to the fact that the inclusive approach is much less sensitive to power in $1/m_c$ corrections which are the main source of corrections discussed above. Indeed, for inclusive widths an expansion in powers of $1/m_b$ exists even for a small mass of final quark.

4 Subleading Form Factors

In difference with $h_{A_1}(1)$ form factors $R_1(1)$, $R_2(1)$ and $R_3(1)$ are not protected against corrections of the first order in $1/m$ at zero recoil. For this reason they can deviate from unity (HQET limit) quite considerably. Theoretical estimates of Ref.⁹ for $R_1(1)$, $R_2(1)$ are

$$R_1(1) \approx 1 + \frac{m_D * -m_c}{2m_c} + \frac{m_B * -m_b}{2m_b} \approx 1.35 ; \quad R_2(1) \approx 0.79 . \quad (18)$$

The theoretical expression for $R_1(1)$ is presented in somewhat simplified form to demonstrate that the quite considerable deviation from the HQET value comes from the $1/m_c$ correction. The predictions (18) are consistent with CLEO experiment¹⁰ where form factors were measured,

$$R_1 = 1.18 \pm 0.30 \pm 0.12 ; \quad R_2 = 0.71 \pm 0.22 \pm 0.07 . \quad (19)$$

If corrections of the first order in $1/m_c$ are so large, then the natural question to ask is about a magnitude of $1/m_c^2$ corrections. The method of sum rules discussed in the previous part for $h_{A_1}(1)$ can be applied to get information about other form factors. Using a set of sum rules from Ref.¹² I came to the following estimate for the $1/m^2$ correction to $R_2(1)$

$$[\Delta R_2(1)]_{1/m^2} \approx \frac{\mu_G^2}{m_c^2 \left(\frac{m_B}{m_{D^*}} - 1 \right)} \left(1 + \frac{m_c}{3m_b} \right) \approx 0.15 \quad (20)$$

The size of the correction is larger than the one of the first order in $1/m$ given by Eq. (18), thus the series in powers of $1/m$ does not look as a convergent expansion. The lesson is that one should be cautious with powers of $1/m_c$.

5 Some Recent Developments

This is a brief mention of a few recent studies relevant to the topic:

- Constraints for form factors following from inclusive sum rules are explored for different semileptonic decays¹³. It results in upper and lower bounds for transitions of B meson to D , D^* , ρ , π , ω , K and
- Inclusive sum rules following from polarization operator of the semileptonic current are used to put a new type of bound for the cross channel (like $B D$) and in this way to the form factors¹⁴.
- Extraction of $|V_{ub}|$ from exclusive modes¹⁵. Using the set of transitions, $B \rightarrow \rho$, $B \rightarrow K^*$, $D \rightarrow \rho$, $D \rightarrow K^*$ it is possible to diminish considerably a model dependence.
- Semileptonic B decays to excited states¹⁶. This interesting development is presented at this conference by Zoltan Ligeti (see his talk).

6 Conclusions

- The theory of semileptonic decays is in a good shape
- The main source of theoretical uncertainties is in the $(1/m_c)^n$ corrections, not in the perturbative ones
- Convergence of exclusive and inclusive approaches

Acknowledgments

I am thankful to Zoltan Ligeti, Mikhail Shifman, Nikolai Uraltsev and Mikhail Voloshin for helpful discussions. This work was supported in part by DOE under the grant number DE-FG02-94ER40823.

References

1. E. Eichten and B. Hill, *Phys. Lett.* **B234** (1990) 511;
H. Georgi, *Phys. Lett.* **B240** (1990) 447.
2. M. Neubert, *Phys. Reports* **245** (1994) 259.
T. Mannel, in: *Schladming 1996, Perturbative and nonperturbative aspects of quantum field theory* 387-428; hep-ph/9606299
3. K. Wilson, *Phys. Rev.* **179** (1969) 1499;
K. Wilson and J. Kogut, *Phys. Reports* **12** (1974) 75.
4. I. Bigi, N. Uraltsev and A. Vainshtein, *Phys. Lett.* **B293** (1992) 430;
(E) B297 (1993) 477;
B. Blok and M. Shifman, *Nucl. Phys.* **B399** (1993) 441; 459.
5. P. Ball and V. Braun, *Phys. Rev.* **D49** (1994) 2472.

6. M. Shifman, N.G. Uraltsev and A. Vainshtein, *Phys.Rev.* **D51** (1995) 2217; (E) **D52** (1995) 3149.
7. A. Czarnecki, K. Melnikov and N. Uraltsev, *Complete $O(\alpha_s^2)$ Corrections to Zero-Recoil Sum Rules for $B \rightarrow D^*$ Transitions*, hep-ph/9706311.
8. I. Bigi, M. Shifman and N. Uraltsev *Aspects Of Heavy Quark Theory*, hep-ph/9703290
9. M. Neubert, *Nucl.Phys.* **B416** (1994) 786.
10. CLEO Collaboration (J.E. Duboscq et al.), *Phys.Rev.Lett.* **76** (1996) 3898.
11. M. Luke, *Phys. Lett.* **B252** (1990) 447. See earlier applications in: M. Voloshin and M. Shifman, *Yad. Fiz.* **47** (1988) 801 [*Sov. J. Nucl. Phys.* **47** (1988) 511].
12. I.I. Bigi, M. Shifman, N.G. Uraltsev and A. Vainshtein, *Phys. Rev.* **D52** (1995) 196.
13. C.G. Boyd and I.Z. Rothstein, *Phys.Lett.* **B395** (1997) 96.
14. C.G. Boyd, B. Grinstein and R. F. Lebed, *Nucl.Phys.* **B461** (1996) 493.
15. Z. Ligeti, I.W. Stewart and M. B. Wise, *Comment on V_{ub} from Exclusive Semileptonic B and D Decays*, hep-ph/9711248.
16. A.K. Leibovich, Z. Ligeti, I.W. Stewart and M. B. Wise, *Semileptonic B decays to excited charmed mesons*, hep-ph/9705467; *Phys.Rev.Lett.* **78** (1997) 3995.